

## Original Article

# Voluntary drinking versus imposed drinking in the methodology of investigations about the *drinking-induced thermoregulatory sweating*

Abdollah Hosseinlou<sup>1</sup>, Saeed Khamnei<sup>2</sup>, Masumeh Zamanlu<sup>3</sup>

<sup>1</sup>Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran; <sup>2</sup>Drug Applied Research Center, Tabriz, Iran; <sup>3</sup>Neurosciences Research Center (NSRC), Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

Received July 6, 2014; Accepted August 16, 2014; Epub October 15, 2014; Published October 30, 2014

**Abstract:** Studies have shown that dehydrated humans or animals in a warm environment begin to sweat within seconds to minutes after drinking. This phenomenon is one of the *drinking-induced thermoregulatory responses*; being investigated from different aspects. Our objective is to show the difference of voluntary drinking and imposed drinking in the methodology of these experiments. Six healthy subjects  $23.7 \pm 0.6$  yr old and  $80.7 \pm 5.7$  kg wt were dehydrated by performing mild exercise (ergometer cycling) in a hot and humid chamber ( $38-40^{\circ}\text{C}$ , 20-28% relative humidity). We incorporated two protocols: after dehydration, subjects were allowed to drink water with 1) imposed volumes of 1, 3, 5 ml/kg and 2) voluntary volumes; on four separate days. The sweating rate was measured on the forehead area before and after drinking. Sweating increased markedly just a few minutes after the onset of drinking. The mean sweat rates of the imposed volumes of 1, 3, 5 ml/Kg were  $0.33 \pm 0.15$ ,  $0.31 \pm 0.17$ ,  $0.47 \pm 0.21$  respectively and for the voluntary volume it was  $0.54 \pm 0.19$ . The mean intake in the voluntary trial was  $6.58 \pm 1.14$  ml/Kg, more than the imposed volume of 5 ml/Kg. The trend of the rate of the sweating response in the imposed trials was distinct from the response in the voluntary trial. Conclusion: There exists a difference between voluntary drinking and imposed drinking in the sweating response that follows rehydration. So it is suggested to use the methods of voluntary drinking in the investigations of this phenomenon, to reveal the natural events that happen in the actual circumstances.

**Keywords:** Drinking-induced sweating, thermoregulatory evaporation, post rehydration, voluntary intake, imposed water consumption


## Introduction

Heat stress happening in ordinary life, in athletic competition, in military endeavors, and even in animals, is the focus for many investigations worldwide, numerous ones with the aim of introducing the physiologic and pathologic phenomena and numerous others with the aim of finding novelties for promoting performances [1-8]. Of such phenomena under research are the drinking-induced thermoregulatory responses. During heat stress together with dehydration, happening naturally together, thermoregulation is partly suppressed to save body fluid and circulation, because most of the mechanisms of thermoregulation including sweating, panting in animals, skin vasodilatation and etc., result in fluid loss and/or circulation deterioration. *Drinking* which means fluid compen-

sation for the body, induces the recovery of thermoregulatory responses, and this is called *drinking induced thermoregulatory responses* [1, 9-23]. One of these responses, the *drinking induced sweating* is discussed methodologically in this paper.

Drinking induced sweating has been studied since 1965 in a study of animal model [24] until now in studies of human athletes [25, 26]. The methodology for its research have components of fluid loss and raised temperature induced by physical activity, hot and humid environment, and occasionally prescription of diuretics; thereafter fluid intake mostly by consumption, and sometimes by IV injection or NG tube. Most of the authors have utilized imposed intake, although in reality voluntary intake occurs and there are significant differences between impo-

## Voluntary drinking in methodology of drinking-induced sweating



**Figure 1.** Sweat rates versus volume of intake in the 4 imposed/voluntary trials.

sed and voluntary drinking. It has been proved that subjects prefer various volumes in various conditions [5, 26-32]. Nevertheless, "imposing" may indicate stress and sympathetic stimulation while autonomic nervous system especially the sympathetic branch is involved in mechanisms of sweating, and this may be a confounding factor. Therefore, there may be advantages in methodologies with voluntary intake. This paper is an attempt to demonstrate plainly the difference of voluntary and imposed drinking in studying the *drinking induced sweating*. We have performed two series of experiments, one with imposed drinking and the other with voluntary drinking [5, 25, 26], and comparison of the resulted sweating responses were evaluated to estimate the confounding effect of imposed drinking.

### Materials and methods

#### Subjects

Six healthy male Taekwondo Athletes mean age  $23.7 \pm 0.6$  years old (range: 22-26), mean weight  $80.7 \pm 5.7$  kg, and mean height  $181 \pm 2$  cm, participated in this study. All subjects were

familiarized with the testing procedures. They were normotensive, nonsmokers, and took no medications. The study was approved by the Research Vice-Chancellor of Tabriz University of Medical Sciences and the Ethics Committee of the University, and written informed consent was obtained from all subjects.

#### Procedure

Pretest instructions included eating a light lunch, refraining from drinking any beverage several hours before the experiment, and no exercise on the day of the experiments. Before each experiment, subjects entered a chamber and rested in the sitting position for 30 min at a thermoneutral temperature ( $28^{\circ}\text{C}$ , 20-28% relative humidity). Body weight was measured with light clothing. After drawing a blood

sample by venopuncture as the first control sample, subjects entered another chamber ( $38-40^{\circ}\text{C}$ , 20-28% relative humidity). They performed a mild exercise (ergometer cycling) by alternating 10 min rest and 20 min exercise periods for 90 min, then exercise continued for the last 30 min period to induce a reduction in total body weight through sweating. Total heat exposure time was 120 min. During heat exposure, subjects were under constant observation for indications of any inability to tolerate the experimental conditions (e.g., elevated heart rate, nausea or confusion).

After the cessation of exercise, subjects dried their body and their forehead, and were then weighed. An indwelling cannula was inserted into a large superficial vein in the forearm to collect free-flowing blood samples. Second control blood sample was drawn through the indwelling cannula. The first control blood sample compares the plasma concentrations of sodium before and after heat exposure while second control blood sample is considered as a control to compare them before and after drinking.

## Voluntary drinking in methodology of drinking-induced sweating

**Table 1.** Dehydration profile, and intake volume of each trial: mean (SD)

Trial type	Weight loss (gr) or sweat loss (ml)	Dehydration Percentage%	Posm before procedure*	Posm after procedure*	Intake (ml)
Imposed volume of 1 ml/kg	2205.2 (471.7)	2.73 (0.28)	302.05 (1.58)	309.05 (2.45)	80.75 (13.90)
Imposed volume of 3 ml/kg	2246.8 (552.1)	2.77 (0.37)	305.55 (2.57)	317.10 (4.41)	242.25 (41.72)
Imposed volume of 5 ml/kg	2125.0 (541.5)	2.62 (0.40)	308.35 (1.58)	312.90 (1.88)	403.75 (69.53)
Voluntary intake	1900.0 (209.8)	2.38 (0.19)	308.00 (2.54)	313.25 (3.62)	531.01 (91.99)

\*(mosmol/kg H<sub>2</sub>O).

Before drinking, sweat rate was measured as control, then subjects were allowed to drink tap water at temperatures of 16°C, at volumes of 1, 3, and 5 ml/Kg of body weight and in voluntary volumes in four separate experiment days. Blood samples were drawn through the indwelling cannula at the start of drinking (0 min) and at 3, 9, 12, and 15 minutes after drinking.

### Measurements

Forehead sweat rate was chosen to represent a localized area of sweating and was measured by the weight gain of a covered filter paper disk (96 cm<sup>2</sup>) placed on the skin over the forehead. The front sides of the disks were covered with waterproof tape to prevent evaporation. The disk was left on the skin each 3 minutes. It was enough to reliably detect weight gain and for saturation of the filter paper. The weight of a filter paper disk was gauged using EK-500 G beam balance, accurate to  $\pm 0.01$  g. Body weight was measured using a Seca beam balance, accurate to  $\pm 100$  g. As a result of experimental conditions, a decrease in weight primarily reflects water loss by thermoregulatory responses, mainly sweating.

Plasma sodium concentration was determined by eppendorf flame photometry (model EFOX 5054, Instrumentation Laboratory). Sodium concentrations were measured as milliequivalents per kilogram of water by correcting for total solids. Because sodium and its associated anions account for about 94 percent of the solute in the extracellular compartment (including plasma), plasma osmolality can be roughly approximated as: Posm = 2.1 \*Plasma sodium concentration [33].

We applied this formula for the estimation of plasma osmolality from plasma sodium concentration.

### Statistics

Data were analyzed by SPSS16. The difference in variables was assessed using paired t test

(one measurement) and repeated measures analysis of variance, ANOVA (multiple measurements). Values of  $P < 0.05$  were considered statistically significant, and all data are presented as means  $\pm$  SD.

### Results

After dehydration procedure, subjects showed a highly significant decrease in their weight ( $P < 0.01$ ), with mean dehydration of 2.63%. Also, their plasma osmolality (Posm) significantly raised from mean of 305.99 before procedure to mean of 313.08 after procedure ( $P < 0.05$ ) and it was statistically similar in the rest of the experiments until 15 minutes after drinking ( $P > 0.05$ ). The weight loss and Posm change did not vary among the trials ( $P > 0.05$ ). The mean intake in the voluntary trial was 6.58 (1.14) ml/Kg. The parameters affected by dehydration, together with the intake of each trial is shown in **Table 1**.

Abrupt increase in sweating occurred 3 minutes after drinking ( $P < 0.05$ ). The sum of all measured sweatings of the subjects was used to calculate the mean sweat rate in each trial, demonstrated versus the volume of intake in **Figure 1**.

### Discussion

Our results show the prominent difference of the sweating response after voluntary intake and imposed intake. After significant dehydration demonstrated by raise in Posm (**Table 1**), subjects voluntarily drank  $6.58 \pm 1.14$  ml/Kg of tap water, more than the imposed volumes of 5 ml/Kg. It has been reported previously that the volume of fluid intake varies due to beverage temperature [5, 16, 26-28, 30, 34-41], taste [28, 31, 32, 40-42], and composition [4, 8, 14, 29, 43-45]. Beverage temperature of nearly 16°C, which equals the temperature of cool tap water, has been introduced as optimum for rehydration and fluid balance [5, 30, 31]. Drinks containing electrolytes, glucose, and fruit taste,

now produced as *sport drinks*, have been shown to induce more intake and more drinking pleasure [4, 8, 17, 28, 29, 31, 32, 38, 39, 41, 44-46]. However, we used plain water which is the most available beverage in daily life. The sweating response, and even the hormonal reactions involved, vary according to the volume of intake [25], so the sweating response of voluntary intake compared to the imposed intake should vary due to the difference in volumes of consumption. The volume of intake in voluntary trial was more than 5 ml/Kg, and the induced sweating response was more than the sweating response of imposed intake of 5 ml/Kg, but the SD bar of voluntary trial in **Figure 1** has fallen quite apart from the trend of imposed trial, apparently lower. This could be better assessed if the imposed volumes were chosen equal to the voluntary volume, with the imposed trial performed after the voluntary trial, however, if the imposed volume is too close to what the subject would prefer, the effect of *imposing* may be masked to some extent. Surely, more studies with different volumes could make this fact more clear. Anyhow, the reason for apparently lower sweating response may be changes in hormonal reactions. During heat stress and rehydration, plasma levels of vasopressin, epinephrine and norepinephrine change, and are dependant to thirst, intake volume and other factors [2, 3, 25, 35, 47-54]. Epinephrine and norepinephrine are sympathetic hormones changing with stress and affecting sweating as an autonomic output [33, 55]. Imposing consumption of a determined volume may burden stress and influence the sympathetic epinephrine and norepinephrine release, confoundingly raising the sweating response. As can be seen (**Figure 1**), the increase in sweating response is sharper in higher volumes, which may indicate more sympathetic confound with imposing higher volume intake.

Ultimately, it can be concluded that imposed drinking may have confounding effects in the methodology of investigations about the drinking-induced sweating, and even for any study about rehydration, and future studies had better consider voluntary intake.

### Acknowledgements

We wish to express our appreciation to our subjects who voluntarily participated in this experiment. This investigation was supported by Drug

Applied Research Center, Tabriz University of Medical Sciences, Tabriz, I. R. Iran.

### Disclosure of conflict of interest

The authors declare that they have no conflict of interest.

**Address correspondence to:** Dr. Saeed Khamnei, Department of Physiology, Islamic Azad University - Tabriz Branch, Manzarieh Street, Tabriz, Postal Code 5166614766, Iran. Tel: 00984113364664; Fax: 00984113364664; E-mail: khamnei\_s@yahoo.com; khamneh@tbzmed.ac.ir

### References

- [1] Libert JP, Candas V, Amoros C, Sagot JC, Lenzi P. Temperature regulation during intermittent exercise with progressive dehydration. *Jpn J Physiol* 1986; 36: 253-66.
- [2] Ramsay DJ, Thrasher TN. Regulation of fluid intake in dogs following water deprivation. *Brain Res Bull* 1991; 27: 495-9.
- [3] Figaro MK, Mack GW. Regulation of fluid intake in dehydrated humans: role of oropharyngeal stimulation. *Am J Physiol* 1997; 272: R1740-6.
- [4] Shirreffs SM, Maughan RJ. Restoration of fluid balance after exercise-induced dehydration: effects of alcohol consumption. *J Appl Physiol* (1985) 1997; 83: 1152-8.
- [5] Khamnei S, Hosseinlou A, Zamanlu M. Water temperature, voluntary drinking and fluid balance in dehydrated taekwondo athletes. *J Sports Sci Med* 2011; 10: 718-24.
- [6] Sawka MN, Francesconi RP, Young AJ, Pandolf KB. Influence of hydration level and body fluids on exercise performance in the heat. *JAMA* 1984; 252: 1165-9.
- [7] Sosa Leon LA. Treatment of exercise-induced dehydration. *Vet Clin North Am Equine Pract* 1998; 14: 159-73.
- [8] Maughan RJ, Shirreffs SM. Recovery from prolonged exercise: restoration of water and electrolyte balance. *J Sports Sci* 1997; 15: 297-303.
- [9] Oara K, Okuda N, Takaba S. Thermoregulatory responses to heat and exercise in Japanese and Caucasians. *Int J Biometeorol* 1975; 19: 99-107.
- [10] Gisolfi CV, Wenger CB. Temperature regulation during exercise: old concepts, new ideas. *Exerc Sport Sci Rev* 1984; 12: 339-72.
- [11] Miyagawa T, Ogawa T, Asayama M, Yamashita Y. [Sweating response to abrupt changes in work load]. *Nihon Seirigaku Zasshi* 1985; 47: 17-24.



## Voluntary drinking in methodology of drinking-induced sweating

- [12] Baker MA. Effects of dehydration and rehydration on thermoregulatory sweating in goats. *J Physiol* 1989; 417: 421-35.
- [13] Baker MA, Turlejska E. Thermal panting in dehydrated dogs: effects of plasma volume expansion and drinking. *Pflugers Arch* 1989; 413: 511-5.
- [14] Millard-Stafford ML, Sparling PB, Roskopf LB, DiCarlo LJ. Carbohydrate-electrolyte replacement improves distance running performance in the heat. *Med Sci Sports Exerc* 1992; 24: 934-40.
- [15] Takamata A, Mack GW, Gillen CM, Jozsi AC, Nadel ER. Osmoregulatory modulation of thermal sweating in humans: reflex effects of drinking. *Am J Physiol* 1995; 268: R414-22.
- [16] Wimer GS, Lamb DR, Sherman WM, Swanson SC. Temperature of ingested water and thermoregulation during moderate-intensity exercise. *Can J Appl Physiol* 1997; 22: 479-93.
- [17] Galloway SD, Maughan RJ. The effects of substrate and fluid provision on thermoregulatory, cardiorespiratory and metabolic responses to prolonged exercise in a cold environment in man. *Exp Physiol* 1998; 83: 419-30.
- [18] Morimoto T, Itoh T. Thermoregulation and body fluid osmolality. *J Basic Clin Physiol Pharmacol* 1998; 9: 51-72.
- [19] Brozmanova A, Jochem J, Javorka K, Zila I, Zwirska-Korczała K. Effects of diuretic-induced hypovolemia/isosmotic dehydration on cardiorespiratory responses to hyperthermia and its physical treatment in rabbits. *Int J Hyperthermia* 2006; 22: 135-47.
- [20] Brozmanova A, Jochem J, Javorka K, Zila I, Zwirska-Korczała K. Diuretic-induced dehydration/hypovolemia inhibits thermal panting in rabbits. *Respir Physiol Neurobiol* 2006; 150: 99-102.
- [21] McKinley MJ, Weissenborn F, Mathai ML. Drinking-induced thermoregulatory panting in rehydrated sheep: influences of oropharyngeal/esophageal signals, core temperature, and thirst satiety. *Am J Physiol Regul Integr Comp Physiol* 2009; 296: R1881-8.
- [22] Lee JY, Wakabayashi H, Wijayanto T, Hashiguchi N, Saat M, Tochihara Y. Ethnic differences in thermoregulatory responses during resting, passive and active heating: application of Werner's adaptation model. *Eur J Appl Physiol* 2011; 111: 2895-905.
- [23] Longhurst JC, Spilker HL, Ordway GA. Cardiovascular reflexes elicited by passive gastric distension in anesthetized cats. *Am J Physiol* 1981; 240: H539-45.
- [24] Senay LC Jr, Christensen ML. Cardiovascular and sweating responses to water ingestion during dehydration. *J Appl Physiol* 1965; 20: 975-9.
- [25] Khamnei S, Hosseini A and Ibrahimi H. The Effect of Volume of Consumed Water on Drinking-Induced Sweating and Plasma Levels of Arginine Vasopressin, Epinephrine and Norepinephrine. *Int J Endocrinol Metab* 2004; 2: 19-28.
- [26] Hosseini A, Khamnei S, Zamanlu M. The effect of water temperature and voluntary drinking on the post rehydration sweating. *Int J Clin Exp Med* 2013; 6: 683-7.
- [27] Zetou E, Giatsis G, Mountaki F, Komninakidou A. Body weight changes and voluntary fluid intakes of beach volleyball players during an official tournament. *J Sci Med Sport* 2008; 11: 139-45.
- [28] Szlyk PC, Sils IV, Francesconi RP, Hubbard RW, Armstrong LE. Effects of water temperature and flavoring on voluntary dehydration in men. *Physiol Behav* 1989; 45: 639-47.
- [29] Maughan RJ, Leiper JB, Shirreffs SM. Factors influencing the restoration of fluid and electrolyte balance after exercise in the heat. *Br J Sports Med* 1997; 31: 175-82.
- [30] Butudom P, Barnes DJ, Davis MW, Nielsen BD, Eberhart SW, Schott HC 2nd. Rehydration fluid temperature affects voluntary drinking in horses dehydrated by furosemide administration and endurance exercise. *Vet J* 2004; 167: 72-80.
- [31] Szlyk PC, Sils IV, Francesconi RP, Hubbard RW, Matthew WT. Variability in intake and dehydration in young men during a simulated desert walk. *Aviat Space Environ Med* 1989; 60: 422-7.
- [32] Hubbard RW, Sandick BL, Matthew WT, Francesconi RP, Sampson JB, Durkot MJ, Maller O, Engell DB. Voluntary dehydration and alliesthesia for water. *J Appl Physiol Respir Environ Exerc Physiol* 1984; 57: 868-73.
- [33] Guyton A and Hall JE. Text book of medical physiology. Philadelphia: Elsevier Saunders; 2011. pp. 358.
- [34] Lee JK, Shirreffs SM, Maughan RJ. Cold drink ingestion improves exercise endurance capacity in the heat. *Med Sci Sports Exerc* 2008; 40: 1637-44.
- [35] Salata RA, Verbalis JG, Robinson AG. Cold water stimulation of oropharyngeal receptors in man inhibits release of vasopressin. *J Clin Endocrinol Metab* 1987; 65: 561-7.
- [36] Galloway SD, Maughan RJ. Effects of ambient temperature on the capacity to perform prolonged cycle exercise in man. *Med Sci Sports Exerc* 1997; 29: 1240-9.
- [37] Lee JK, Shirreffs SM. The influence of drink temperature on thermoregulatory responses during prolonged exercise in a moderate environment. *J Sports Sci* 2007; 25: 975-85.

## Voluntary drinking in methodology of drinking-induced sweating

- [38] Lee JK, Maughan RJ, Shirreffs SM. The influence of serial feeding of drinks at different temperatures on thermoregulatory responses during cycling. *J Sports Sci* 2008; 26: 583-90.
- [39] Sandick BL, Engell DB, Maller O. Perception of drinking water temperature and effects for humans after exercise. *Physiol Behav* 1984; 32: 851-5.
- [40] McBurney DH, Collings VB, Glanz LM. Temperature dependence of human taste responses. *Physiol Behav* 1973; 11: 89-94.
- [41] Boulze D, Montastruc P, Cabanac M. Water intake, pleasure and water temperature in humans. *Physiol Behav* 1983; 30: 97-102.
- [42] McBurney DH, Bartoshuk LM. Interactions between stimuli with different taste qualities. *Physiol Behav* 1973; 10: 1101-6.
- [43] Hickey MS, Costill DL, Trappe SW. Drinking behavior and exercise-thermal stress: role of drink carbonation. *Int J Sport Nutr* 1994; 4: 8-21.
- [44] Lee JK, Maughan RJ, Shirreffs SM, Watson P. Effects of milk ingestion on prolonged exercise capacity in young, healthy men. *Nutrition* 2008; 24: 340-7.
- [45] Burke LM. Nutritional needs for exercise in the heat. *Comp Biochem Physiol A Mol Integr Physiol* 2001; 128: 735-48.
- [46] Galloway SD, Maughan RJ. The effects of substrate and fluid provision on thermoregulatory and metabolic responses to prolonged exercise in a hot environment. *J Sports Sci* 2000; 18: 339-51.
- [47] Geelen G, Greenleaf JE, Keil LC. Drinking-induced plasma vasopressin and norepinephrine changes in dehydrated humans. *J Clin Endocrinol Metab* 1996; 81: 2131-5.
- [48] Ramsay DJ. The importance of thirst in maintenance of fluid balance. *Baillieres Clin Endocrinol Metab* 1989; 3: 371-91.
- [49] Geelen G, Keil LC, Kravik SE, Wade CE, Thrasher TN, Barnes PR, Pyka G, Nesvig C, Greenleaf JE. Inhibition of plasma vasopressin after drinking in dehydrated humans. *Am J Physiol* 1984; 247: R968-71.
- [50] Appelgren BH, Thrasher TN, Keil LC, Ramsay DJ. Mechanism of drinking-induced inhibition of vasopressin secretion in dehydrated dogs. *Am J Physiol* 1991; 261: R1226-33.
- [51] Thrasher TN, Nistal-Herrera JF, Keil LC, Ramsay DJ. Satiety and inhibition of vasopressin secretion after drinking in dehydrated dogs. *Am J Physiol* 1981; 240: E394-401.
- [52] Lee MC, Thrasher TN, Ramsay DJ. Is angiotensin essential in drinking induced by water deprivation and caval ligation? *Am J Physiol* 1981; 240: R75-80.
- [53] Ramsay DJ, Rolls BJ, Wood RJ. Thirst following water deprivation in dogs. *Am J Physiol* 1977; 232: R93-100.
- [54] Engell DB, Maller O, Sawka MN, Francesconi RN, Drolet L, Young AJ. Thirst and fluid intake following graded hypohydration levels in humans. *Physiol Behav* 1987; 40: 229-36.
- [55] Shirreffs SM, Maughan RJ. Whole body sweat collection in humans: an improved method with preliminary data on electrolyte content. *J Appl Physiol* (1985) 1997; 82: 336-41.