

Case Report

A subcutaneous follicular dendritic cell sarcoma occurring in the cheek: case report

Yohei Kawasaki¹, Teruyuki Sato¹, Yuko Hiroshima², Hiroshi Nanjo², Maya Suzuki³, Yasufumi Omori³, Takechiyo Yamada¹

¹Department of Otorhinolaryngology, Head and Neck Surgery, Akita University Graduate School of Medicine, Akita, Japan; ²Division of Clinical Pathology, Akita University Hospital, Akita, Japan; ³Department of Molecular and Tumor Pathology, Akita University Graduate School of Medicine, Akita, Japan

Received March 14, 2017; Accepted December 2, 2017; Epub February 15, 2018; Published February 28, 2018

Abstract: We report a subcutaneous follicular dendritic cell sarcoma (FDSC) in the cheek. Follicular dendritic cell sarcoma is a rare neoplasm. Here we report the case of an 81-year-old woman presenting with the chief complaint of swelling of the left cheek. We could only diagnose granulation tissue based on a core-needle biopsy; thus, we decided to excise the tissue for definitive diagnosis. At first, we considered a B-cell lymphoma, but the diagnosis of FDSC was made following immunostaining of FDC markers. Tumor resection was performed without any other treatments such as chemoradiotherapy. No recurrence has so far been detected for three years. In the head and neck, FDSC occurs with a predilection for the tonsils. This appears to be the first report of a case of FDSC occurring subcutaneously in the cheek. Currently, there are several options for treatment and can only be standardized by accumulating experience with future cases.

Keywords: Follicular dendritic cell sarcoma, head and neck, subcutaneous, immunohistochemical staining

Introduction

Follicular dendritic cells (FDC) are antigen-presenting cells found in lymphoid follicles that help maintain the lymph-node structure. FDC sarcomas (FDSC) are classified as histiocytic and dendritic cell neoplasms by the World Health Organization and are believed to originate from FDC. Although they mainly occur in lymph nodes of the neck and the abdominal cavity, majority of the cases of the head and neck FDSC arise in the tonsils (**Table 1**). We report a case in which a subcutaneous FDSC developed in the cheek.

Case report

An 81-year-old woman presented with swelling of the left cheek. Blood tests revealed no abnormalities; T1-weighted magnetic resonance imaging (MRI) revealed low signals and T2-weighted MRI revealed a mixture of low and high signals (**Figure 1**). A firm and partially mobile tumor was palpable. The tumor was diagnosed following a core-needle biopsy, but

because a definitive diagnosis could not be made, tumor excision was planned.

During the surgery, an incision under the canine confirmed the presence of the tumor, and because it surrounded the infraorbital nerve, it appeared to be an infraorbital nerve schwannoma. We removed the entire infraorbital nerve, including the surrounding tissues. There was no tumor infiltration in the surrounding tissues, and the tumor was dissectible from the surrounding tissues (**Figure 2**).

Histopathology

Immunohistochemical staining revealed numerous B lymphocytes expressing CD20 and CD10 with few atypical cells in addition to CD3⁺ and CD4⁺ T lymphocytes. We initially considered low-grade B-cell lymphoma [1]. However, HE staining revealed a solid tumor with some whorl formations comprising spindle-shaped and ovoid cells. Immunohistochemical staining was positive for 4 FDC markers, including CD21, CD23, CD35, and CAN.42, and CD79a, and, on the other hand, negative for S-100, α -SMA,

Follicular dendritic cell sarcoma in the cheek

Table 1. Head and neck FDSC reported in literatures [3-5]

Site	Number of cases
Tonsil	62
Parapharyngeal space	4
Soft palate	2
Nasopharynx	6
Hypopharynx	1
Cheek (present case)	1




Figure 1. MRI imaging of the tumor (arrow): MRI showing low-intensity signals in T1-weighted (A and C) and mixture of low-intensity and high-intensity signals in T2-weighted images (B and D).

CD3, EGFR, and ALK-1 (Figure 3). In addition, EBER-1 RNA was not expressed as revealed by *in situ* hybridization. The diagnosis was subcutaneous FDSC in the cheek. The MIB-1 index was 70%, exclusive of non-tumor cells, in hot spots. Tumor resection was performed without any other treatments such as chemoradiotherapy. No recurrence has so far been detected for three years.

Discussion

FDSC is a very rare tumor believed to originate in FDC and was first reported in 1986 by Mondo et al [2]. Majority of the cases show the pres-

ence of such tumors in the lymph nodes, with reports of approximately 80 previous cases occurring in the head and neck region, with a predilection for the tonsils [3-5]. FDSC has been reported in a wide range of patient ages and equally in both sexes.

The etiology of the disease is unknown but may be associated with hepatitis B virus infection or Cattleman's disease [6]. This patient tested negative for Epstein-Barr virus-encoded small RNAs by *in situ* hybridization. Patients are usually negative for CD79a, but our patient was positive for it; in addition, some patients are reported to be negative for UCHL1 (CD45RO) but some other are reported to be positive. Thus, FDSC can have a variety of immunological characteristics. In one report, 26 of 46 cases of FDSC in the head and neck region were initially misdiagnosed; accurate diagnoses were reached only after reassessment. Immunohistochemical staining has shown positive staining for FDC markers including CD21, CD23, CD35, Ki-M4p, CAN.42, and D2-40 [7].

FDSC is a low-grade to intermediate-grade tumor [6] with distant metastasis in 16% of cases originating in the head and neck region and tumor-specific death in 10%. The published disease-free survival rate is 66.2% [8]. In tumor grade classification for estimating prognosis, Ki-67 labeling index <10% is considered low grade and $\geq 10\%$ is considered as high grade [8]. Treatment varies and includes surgery alone, combination therapy with surgery and radiotherapy or chemotherapy, or chemoradiotherapy [9]; in addition, no consensus has been reached regarding the preferred treatment. Patients with recurrence or tumor progression have been treated with cyclophosphamide, doxorubicin, vincristine, and prednisone, which are used in treating




Figure 2. Macroscopic findings: A. Surgical appearance. B. The surface of the excised tissue was smooth. C. The cross-sectional surface of the tumor was grayish white.




Figure 3. Microscopic findings: A (original magnification, $\times 100$) and B ($\times 400$): Whorl formations of tumor cells with unclear cell borders and growing as a sheet. C: CD21-positive immunostaining of the tumor cell cytoplasm ($\times 100$). D ($\times 100$): MIB-1 index was 70%, exclusive of non-tumor cells, in hot spots ($\times 100$).

lymphoma [10-12]. In this patient, the MIB-1 index was 70%; hence, the tumor was believed to be high grade; however, given the advanced age of the patient and considering her overall health condition, additional treatment such as radiotherapy or chemotherapy was not provided. To our knowledge, this is the first report of FDCS occurring in the cheek. Because it is such a rare tumor, it is necessary to compile case reports from around the world to establish a uniform treatment strategy.

Conclusion

To our knowledge, this unusual case is probably the first reported one of subcutaneous FDCS occurring in the cheek. Given that the etiology and preferred treatment are currently not established, additional experiences are needed before recommendations can be made.

Disclosure of conflict of interest

None.

Address correspondence to: Yohei Kawasaki, Department of Otorhinolaryngology, Head and Neck Surgery, Akita University Graduate School of Medicine, Akita, Japan. Tel: +81 18-884-6171; Fax: +81 18-884-2622; E-mail: kawa0807@med.akita-u.ac.jp

References

- [1] Cordes C, Tiemann M, Tiemann K, Knappe D, Hoffmann M and Gottschlich S. Epstein-Barr virus associated diffuse large B-cell lymphoma of the hypopharynx. *B-ENT* 2011; 7: 43-46.
- [2] Mondo L, Warnke R and Rosai J. A primary lymph node malignancy with features suggestive of dendritic reticulum cell differentiation, a report of 4 cases. *Am J Pathol* 1986; 122: 562-572.
- [3] Hu T, Wang X, Yu C, Yan J, Zang X, Li L, Li X, Wu J, Ma W, Li W, Wang G, Zhao W, Zhang D and Zhang M. Follicular dendritic cell sarcoma of the pharyngeal region. *Oncol Lett* 2013; 5: 1467-1476.
- [4] Amirtham U, Manohar V, Kamath MP, Srinivasamurthy PC, Chenagiriappa LK, Shenoy AM, Renuka PK, Kumar RV. Clinicopathological profile and outcomes of follicular dendritic cell sarcoma of the head and neck region-A study of 10 cases with literature review. *J Clin Diag Res* 2016; 10: XC08-XC11.
- [5] Kulkarni MP, Monin YA, Deshmukh BD, Sulhyan KR. Extranodal follicular dendritic sarcoma

Follicular dendritic cell sarcoma in the cheek

- involving tonsil. *Malaysian J Pathol* 2015; 37: 293-299.
- [6] Chan JK, Pileri SA, Delsol G, Fletcher CD, Weiss LM and Grogg KL. Follicular dendritic cell sarcoma, WHO Classification of tumors of haematopoietic and lymphoid tissues in: Swerdlow S, Campo E, Jaffe E, Pileri S, Stein H, Thiele J and James W, editors. 4th edition. Vardiman Lyon; IARC; 2008; pp; 363-364.
- [7] Xie Q, Chen L, Fu K, Harter J, Young KH, Sunkara J, Novak D, Villanueva-Siles E and Ratech H. Podpulanin (D2-40): a new immunohistochemical marker for reactive follicular dendritic cell and follicular dendritic cell sarcoma. *Int J Clin Exp Pathol* 2008; 1: 276-284.
- [8] Chan JK, Fletcher CD, Nayler SJ and Cooper K. Follicular dendritic cell sarcoma; clinicopathologic analysis of 17 case suggesting a malignant potential higher than currently recognized. *Cancer* 1997; 179: 294-313.
- [9] Masunaga A, Nakamura H, Kanata T, Furubayashi T, Kanayama Y, Yamada A, Shiroko Y and Itoyama S. Follicular dendritic cell tumor with histiocytic characteristics and fibroblastic antigen. *Pathol Int* 1997; 47: 707-712.
- [10] Li L, Shi YH, Guo ZJ, Qiu T, Guo L, Yang HY, Zhang X, Zhao XM and Su Q. Clinicopathological features and prognosis assessment of extranodal follicular dendritic cell sarcoma. *World J Gastroenterol* 2010; 16: 2504-2519.
- [11] Cyriac S, Praveenkumar D, Majhi U and Sagar TG. Follicular dendritic cell sarcoma of the neck with an aggressive and fatal course. *J Cancer Res Ther* 2010; 6: 114-116.
- [12] Pisani F, Marino M, Sentinelli S and Petti MC. Follicular dendritic cell sarcoma of the neck report of a case treated by surgical excision and COP plus (PEG)-liposomal doxorubicin. *J Exp Clin Cancer Res* 2008; 27: 33.